

A Variant Approach to the Overlap Action

Thomas DeGrand

(MILC Collaboration)

Department of Physics, University of Colorado, Boulder, CO 80309 USA

(February 1, 2008)

I describe an implementation of the overlap action, which is built from an action which is itself an approximate overlap action. It appears to be about a factor of 15-20 less expensive to use, than the usual overlap action with the Wilson fermion action as its kernel. Ingredients include a fat link to suppress coupling to dislocations and a free field action with a spectrum which resembles an overlap; much of the gain comes from the use of eigenmodes of the approximate action to begin the overlap calculation. As a physics example, I compute the quark condensate in finite volume in the quenched approximation.

I. INTRODUCTION

There has been a great deal of interest recently in lattice fermion actions which implement an exact chiral symmetry without doubling [1]. The one explicit realization of such an action is the overlap action of Neuberger [2]. It obeys the simplest version of the Ginsparg-Wilson [3] (G-W) relation. All published studies of the overlap action in four dimensions [4–12] use the Wilson fermion action as their starting point. These actions are apparently very expensive to simulate. However, it is a common expectation that a better starting action would simplify the evaluation of an overlap action, if many fewer steps of iteration would compensate for the increased cost per step of the more complicated action. That expectation is realized by an action (or family of actions) I describe here. With it, some physics calculations using an overlap action in quenched approximations can be done without recourse to supercomputers. Ingredients in my scheme include a fat link to suppress coupling to dislocations and a free field action with a spectrum which resembles an overlap; much of the gain comes from the use of eigenmodes of the approximate action to begin the calculation of overlap eigenmodes.

These techniques are obviously inspired by the fixed point action program for constructing classically perfect fermion actions [13]. The best action I have found is, however, not, as far as I know, a fixed point action of any renormalization group transformation.

Versions of these ideas have been presented many times in two dimensional models [14].

After setting some conventions in Sec. II, I will describe the candidate actions and their tests in Sec. III. I will then present in Sec. IV a calculation of the quark condensate in finite volume with one of the new actions, along the lines of the calculation of Ref. [11].

II. NOTATION AND CONVENTIONS

I will call a generic lattice Dirac operator d and the overlap Dirac operator D . The eigenvalues of the simplest implementation of a G-W action lie on a circle of radius r_0 and the (massless) overlap Dirac operator is

$$D(0) = r_0(1 + \frac{z}{\sqrt{z^\dagger z}}) \quad (1)$$

where $z = d(-r_0)/r_0 = (d - r_0)/r_0$ and $d(m) = d + m$ is the massive Dirac operator for mass m (i.e. r_0 is equivalent to a negative mass term.) The overall multiplicative factor of r_0 is a useful convention; when the Dirac operator d is thought of as “small” and Eqn. 1 is expanded for small d , $D \simeq d$. Apart from this overall factor of r_0 , my conventions are those of Ref. [7].

The Hermitian Dirac operator for mass m is defined as $h(m) = \gamma_5 d(m)$ and the overlap Hermitian Dirac operator is denoted as $H(m)$. Specifically,

$$H(0) = r_0(\gamma_5 + \epsilon(h(-r_0))) \quad (2)$$

where $\epsilon(x)$ is the step function, $\epsilon = -1$ if $x < 0$, $\epsilon = 1$ if $x > 0$. I will refer to the argument of the step function as the “kernel” of the overlap.

It is convenient to define the massive overlap Dirac operator as

$$D(m) = (1 - \frac{m}{2r_0})D(0) + m \quad (3)$$

and then the squared massive Hermitian Dirac operator is

$$H(m)^2 = m^2 + (1 - \frac{m^2}{4r_0^2})H(0)^2. \quad (4)$$

The zero eigenvalue eigenmodes of $H(0)$ are chiral, with $\langle \phi | \gamma_5 | \phi \rangle = \pm 1$, and the nonzero eigenvalue eigenmodes of $H(0)$ come in pairs of equal and opposite eigenvalues. For these modes, $\langle \phi_\lambda | \gamma_5 | \phi_\lambda \rangle = \langle \phi_\lambda | H(0) | \phi_\lambda \rangle / (2r_0) = \lambda / (2r_0)$.

In a background gauge field carrying a topological charge Q , $D(0)$ (and $H(0)$) will have Q pairs of real eigenmodes with eigenvalues 0 and $2r_0$. In computing propagators (for example for $\langle \psi \psi \rangle$), it is convenient to clip out the eigenmode with real eigenvalue $2r_0$, and to define the subtracted propagator as

$$\tilde{D}(m)^{-1} = \frac{1}{1 - \frac{m}{2r_0}} [D(m)^{-1} - \frac{1}{2r_0}]. \quad (5)$$

III. THE OVERLAP WITH A BETTER KERNEL

A. Ingredients

A good d or h for use as a kernel in Eq. 1 or 2 should already “look like” D or H . This means that the eigenvalues of its eigenmodes should lie (approximately) on a circle and its low lying eigenmodes should be approximately chiral—in a spectroscopy calculation, the additive renormalization of the bare quark mass, as measured, for example, through the variation of the pion mass with bare quark mass, should be small. Most implementations of the overlap action in the literature use the Wilson action, which does not satisfy either of these criteria: the eigenvalues of the free Wilson action sit like beads on a string around a set of four circular arcs (with real parts of their eigenvalues ranging from 0 to 8), and the additive mass renormalization of the interacting theory is $\Delta am_q \simeq 1$ for simulations at lattice spacings near 0.15 fm. Use of the clover action instead of the Wilson action improves the chiral properties but does nothing for its (free field) eigenmode spectrum.

Improving the action involves two ingredients.

First, to improve the chiral properties of the action, thin links are replaced by fat links. Actions with fat links are already quite chiral as shown by their small mass renormalization and $Z_A \simeq 1$ [15,16]. In this work I have studied fat link actions with two blockings: The first uses APE-blocking [17]:

$$\begin{aligned} V_\mu^{(n)}(x) = & (1 - \alpha)V_\mu^{(n-1)}(x) \\ & + \alpha/6 \sum_{\nu \neq \mu} (V_\nu^{(n-1)}(x)V_\mu^{(n-1)}(x + \hat{\nu})V_\nu^{(n-1)}(x + \hat{\mu})^\dagger \\ & + V_\nu^{(n-1)}(x - \hat{\nu})^\dagger V_\mu^{(n-1)}(x - \hat{\nu})V_\nu^{(n-1)}(x - \hat{\nu} + \hat{\mu})), \end{aligned} \quad (6)$$

with $V_\mu^{(n)}(x)$ projected back onto $SU(3)$ after each step, and $V_\mu^{(0)}(n) = U_\mu(n)$ the original link variable. I have mostly studied a large amount of fattening, $\alpha = 0.45$, $N = 10$, but have looked at the smaller value $\alpha = 0.25$, $N = 7$. The second blocking is a “hypercubic” blocking devised by A. Hasenfratz [18]. The fat links of this action are confined to a hypercube, so this action is more local than the APE-blocked actions. The mean plaquette at $\beta = 5.9$ for (0.45,10) APE blocking is about 2.98, for (0.25,7) APE blocking it is 2.88, and for the hypercubic blocking, 2.84.

I believe that what is important here is that the fat gauge links decouple the fermions from short distance structure in the gauge field, not that the links are fattened in a particular way.

The eigenvalues of a GW action lie on a circle. I determine the best d by taking a free field test action and varying its parameterization to optimize its eigenvalue spectrum (in the least-squares sense) for circularity, for some r_0 . It was convenient to let the parameter r_0 also be a free parameter. The simplest extension of a nearest neighbor action which can have its eigenvalues lying approximately on a circle is “planar:” it has scalar and vector couplings $S = \sum_{x,r} \bar{\psi}(x)(\lambda(r) + i\gamma_\mu \rho_\mu(r))\psi(x+r)$ for r connecting nearest neighbors ($\vec{r} = \pm \hat{\mu}$; $\lambda = \lambda_1 = -0.170$, $\rho_\mu^{(1)} = -0.177$)

and diagonal neighbors ($\vec{r} = \pm\hat{\mu} \pm \hat{\nu}$, $\nu \neq \mu$; $\lambda = \lambda_2 = -0.061$, $\rho_\mu^{(2)} = \rho_\nu^{(2)} = -0.0538$. The constraint $\lambda(r=0) = -8\lambda_1 - 24\lambda_2$ enforces masslessness on the spectrum, and $-1 = 2\rho_\mu^{(1)} + 12\rho_\mu^{(2)}$ normalizes the action to $-\bar{\psi}i\gamma_\mu\partial_\mu\psi$ in the naive continuum limit. The corresponding value of r_0 is 1.6. A plot of the free field eigenvalues on a finite lattice is shown in Fig. 1. Presumably the places where its eigenvalues are purely real (but not zero) correspond in the continuum limit to theories with various numbers of massless free fermions, exactly as for the Wilson action.




FIG. 1. Eigenmode spectrum for the free field planar action on a finite lattice. Only the positive imaginary part modes are shown.

The action also includes a clover term, with its coefficient set to the tree-level value appropriate to this action of $C_{SW} = 1.029941$. Thus d has no $O(a)$ artifacts. All the links, including the ones in the clover term, are replaced by fat links. The gauge connections to the diagonal neighbors are the average of the two shortest path connections. The cost of the action is about 6.5 times that of the usual clover action.

While it is probably not germane to the present discussion, the dispersion relation of this action is improved compared to that of clover or Wilson actions.

No claims are made for uniqueness—or even optimality. The optimizations do not have to be done with enormous precision since the overlap formula itself deforms almost any kernel action into a chiral action. The amount of fattening is also a free parameter, and there will almost certainly be tradeoffs between ease of implementing the overlap and the desire that the action show good scaling behavior. At large levels of fattening, there appears to be little renormalization of the parameters of a free action, and so little tuning is required to produce actions which behave well in simulations. Thus, the overlap parameter r_0 will be kept at its free field value of 1.6.

B. Testing actions

In what follows I will focus on comparisons of the overlap action with the Wilson action as a kernel, the “Wilson overlap” and with the fat link planar action as a kernel, or “planar overlap.” I have looked at six actions in all. Most tests involve the Wilson overlap or the (0.45,10) APE-blocked planar overlap. I have briefly investigated an overlap action with the (0.45,10) APE-blocked fat link clover action as a kernel. I have also investigated planar overlaps with (0.45,10) APE-blocked links and with hypercube-blocked links. Finally, I tested the “Gaussian” action, which has been proposed as a candidate approximate FP action [16].

The step function is approximated by either the polar formula introduced by Neuberger

$$\epsilon(z) \rightarrow \epsilon_N(z) = z \frac{1}{N} \sum_{j=1}^N \frac{1}{c_j z^2 + s_j} \quad (7)$$

($c_j = \cos^2(\pi(j + 1/2)/(2N))$, $s_j = \sin^2(\pi(j + 1/2)/(2N))$), or by a fourteenth order Remes polynomial, following the work of Edwards, Heller and Narayanan [7]. In practice, the polar formula works very robustly for the planar overlap, when I rescale $h(-r_0)$ to $h(-r_0)/r_0$ in Eq. 7 and take $N = 6$ to 10. The Wilson overlap is much more delicate and I have used the Remes approximation exclusively for it. I rescaled the operator $h(-r_0)$ by a factor of 1/2.5 to map its eigenvalues into a range where the Remes algorithm has small errors. The inverses of the terms in the sum of Eq. 7 are found using a multi-mass conjugate gradient routine [19].

Let’s begin by looking at the actions. The range of the action is computed using a variation of the calculation of Hernández, Jansen, and Lüscher [11]: Shown in Figs. 2 and 3 is a comparison of Wilson and planar overlap $\sqrt{|D(0)\chi|^2}$ for a delta-function source χ at the origin, as a function of distance $r = \sqrt{x_\mu^2}$, for the free field action as well as on a set of $\beta = 5.9$ 8^4 configurations. The two actions show similar exponential falloff with distance. Of course, these pictures do not show anything about the locality of the gauge connections in the actions, just the fermions. In perturbation theory [20] the fat link action has a cloud of glue of size $\sqrt{\alpha N/3}$ convoluted over every fermion offset.




FIG. 2. Range of the Wilson overlap action, in free field theory and on a set of $\beta = 5.9$ 8^4 lattices.




FIG. 3. Range of the (0.45,10) APE-blocked planar overlap action, in free field theory and on a set of $\beta = 5.9$ 8^4 lattices.

It might be relevant that the plaquette for the fat link action has a value $\text{Tr}(1 - U_p)/3 \simeq 1 - 2.97/3 = 0.01$, which should be sufficient for the argument of Hernández, Jansen, and Lüscher [5] to insure the locality of the overlap action.

All my tests of the overlap begin with finding the eigenmodes of H with the smallest eigenmodes (actually the eigenmodes of H diagonalized in a basis which is composed of the smallest eigenmodes of H^2). Low lying eigenmodes of $h(m)$ and $H(0)$ are found using an adaption of a conjugate gradient algorithm of Bunk et. al. and Kalkreuter and Simma [21]; I modified a code originally written for staggered fermions by K. Orginos.

I have implemented most of the standard tricks for efficient evaluation of eigenmodes of H . I compute a set of the N_0 low lying eigenmodes of $h(-r_0)$ (typically $N_0 = 10 - 20$), and project them out during the calculation of the operator $\epsilon(h(-r_0))\chi$.

A first sign that the fat link will make the overlap better behaved comes from looking at the eigenmodes of the kernel function $h(-r_0)$. Fig. 4 shows a histogram of the ten smallest eigenmodes of $h(-r_0)$ reconstructed from the ten smallest eigenmodes of $h(-r_0)^2$ on a set of ten 8^4 $\beta = 5.9$ configurations. The fat link actions have many fewer small eigenmodes. Eigenmodes near zero are associated with dislocations, gauge configurations on which topological objects are about to disappear. It is difficult for approximations to the step function to process these modes. The absence of these modes in the spectrum of $h(-r_0)$ is why the polar form of ϵ_N can perform well, even for small N . There is a lot of discussion in the literature (see for example Ref. [22]) about improving overlap or domain wall fermions by using gauge actions which have fewer dislocations than the Wilson gauge action, but a fermion action which could not see dislocation would work just as well. A fat link action does just that. While the fat link clover action is as much as an improvement as the planar action, it has a much higher conditioning number than the planar action. This is reflected in the amount of work needed to find the eigenvalues (and will also affect the number of inner conjugate

gradient steps needed to evaluate the step function): the average number of Rayleigh iterations needed to collect the ten lowest eigenmodes is 3600 for the Wilson action, 5100 for the fat link clover action, and 1900 for the (fat link) planar action.




FIG. 4. Spectrum of $h(-r_0)$ for the ten smallest eigenvalues of $h(-r_0)^2$ for (a) the Wilson action ($r_0 = 1.65$), (b) the fat link planar action ($r_0 = 1.6$), and the fat link clover action ($r_0 = 1.0$), on $10 \cdot 8^4$ $\beta = 5.9$ configurations.

Because $[H(0)^2, \gamma_5] = 0$, eigenmodes of $H(0)^2$ can be simultaneously eigenmodes of γ_5 . All eigenmodes are constructed in a basis of chiral eigenfunctions. In that basis, eigenmodes of $H(0)^2$ are expected to be doubly degenerate. This degeneracy can be used to test algorithms. A low order polar approximation to the step function does a poor job of producing degenerate pairs of eigenmodes of $H(0)^2$ for the Wilson overlap and at this level of test it must be discarded.

The calculation of the condensate in Sec. IV requires the quark propagator $D(m)^{-1}$, with the contribution from chiral modes removed. I do this by taking the set of eigenmodes of h and finding the two lowest eigenmodes $H(0)^2$ in a basis containing one state of each chirality. The chirality of the highest state identifies the chiral sector which has no zero modes. I then perform the inversion of $H^2 = D^\dagger D$ in this sector. This calculation is also accelerated by finding and projecting out some large number ($O(10)$) of eigenmodes of $H(0)^2$ during this inversion.

A trick which I have not noticed to have been emphasized in the literature is actually the source of much of the gain in efficiency of the planar overlap. This is to begin the calculation of eigenmodes of H by first finding a set of low lying eigenmodes of $h(m)$ for some potentially useful m , and beginning the calculation for H using the eigenmodes of h (rather than, say, beginning with a set of random vectors). The combination of the small number of small eigenmodes

of $h(-r_0)$ for the fat link action and the use of good trial functions makes the planar overlap quite efficient compared to the Wilson overlap.

Most of my tests are on a set of 8^4 $\beta = 5.9$ lattices. Let us consider a set of examples which illustrate the differences. In the planar overlap I take the $N = 10$ polar approximation to the step function, scale the argument by $r_0 = 1.6$, and project out $N_0 = 10$ eigenmodes of $h(-r_0)$ in the step function. The Wilson overlap uses the fourteenth-order Remes approximation, scales its argument by $1/2.5$, and projects 20 eigenvalues of $h(-r_0)$. In both cases I wish to find the lowest two eigenmodes of $H(0)^2$, regardless of chirality, and I begin by finding and utilizing the four smallest eigenmodes of $h(0)$. I investigate a configuration which happens to have topological charge $Q = 1$. The planar overlap calculation needs 109 Rayleigh iterations and 2529 inner conjugate gradient steps to find the two lowest modes (at a cost of 6.5 times the equivalent Wilson action step). The Wilson overlap needs 1220 Rayleigh iterations, and 86,723 inner conjugate gradient steps, to reach the same level of accuracy (about 10^{-5} for the eigenvalues). Had we wanted the ten smallest eigenmodes of $H(0)^2$, using 20 trial eigenmodes of $h(0)^2$, the cost would be 306 Rayleigh iterations and 8189 inner conjugate gradient steps for the planar overlap, or about 26 inner steps per Rayleigh iteration, while the Wilson overlap uses 5820 Rayleigh iterations and a million inner conjugate gradient steps (about 170 inner steps per iteration). This is a savings in computer time of about a factor of eighteen for the planar overlap compared to the Wilson overlap. On other configurations, the number of inner CG's per Rayleigh step for the planar overlap ranges between 20 and 30, and the number of Rayleigh iterations ranges from 300 to 500.

The difference in the number of inner conjugate gradient steps is due to the fact that the Wilson $h(-r_0)$ has many more small eigenvalues than the fat link planar action. The difference in the number of Rayleigh iterations arises because the initial trial vectors, eigenmodes of $h(0)$, are close to being eigenmodes of $H(0)$. I illustrate this with a scatter plot of the change in an eigenmode, plotted as a function of the value of the overlap eigenmode, in Fig. 5. A graph of $\langle \gamma_5 \rangle$ vs. eigenmode for this action, at various stages of the overlap calculation, is shown in Fig. 6. It is also useful to look at the scatter of chirality vs. eigenvalue for several trial actions which might be candidates for an overlap kernel. This is shown in Fig. 7. I show $\langle \gamma_5 \rangle$ vs. eigenmode for the Wilson action, non-perturbative thin link clover action (with $C_{SW} = 1.85$), fat link planar action, and fat link Gaussian action. From this picture, one would suspect that the thin link actions would not provide good trial wave functions for the overlap.




FIG. 5. Differences between exact eigenvalues of $H(0)$ and (a) eigenvalues of the planar $h(0)$ and (b) eigenvalues of $H(0)$ produced by diagonalizing $H(0)$ in the lowest 20 eigenvalue eigenmodes of $h(0)^2$, as a function of overlap eigenvalue.




FIG. 6. $\langle \gamma_5 \rangle$ vs. eigenvalue for the planar overlap: (a) 20 smallest eigenvalues of $h(0)$, (b) 20 smallest eigenvalues of $H(0)$ from a diagonalization of the basis of (a), (c) 20 smallest eigenvalues of $H(0)$ from a full calculation, all on a set of ten $\beta = 5.9$ 8^4 lattices.




FIG. 7. $\langle \gamma_5 \rangle$ vs. eigenvalue for several $h(0)$'s. (a) thin link Wilson action, (b) non-perturbative thin link clover action, (c) Gaussian action, (d) planar action.

I have tested another fat link action, a planar action with 7 APE-blocking steps and $\alpha = 0.25$. I kept $r_0 = 1.6$ since tests of the fat link clover action with this level of fattening at $\beta = 5.9$ also showed little additive mass renormalization. The number of small eigenvalues of $h(-r_0)$ increases slightly compared to the (0.45,10) case, and the number of inner CG steps grows, correspondingly, to about 30. But the eigenmodes of $h(0)$ do not seem to be as good a trial basis as they were for the fatter link action, and the number of Rayleigh iterations grows from about 10,000 to 12,000-15,000 for the same calculation as was done above. A set of pictures showing relevant information for this action is shown in Fig. 8. This action does not see some of the instantons that the (0.45,10) fattening saw. That is reflected in the scatter of the eigenvalues.




FIG. 8. $\langle \gamma_5 \rangle$ vs. eigenvalue for the planar action and its overlap with (7,0,25) fattening: (a) 20 smallest eigenvalues of $h(0)$, (b) 20 smallest eigenvalues of $H(0)$ from a diagonalization of the basis of (a), (c) histogram of ten smallest eigenmodes of $h(-r_0)$, on $\beta = 5.9 \cdot 8^4$ configurations.

One might hope that a better trial function would help. A simple way of changing the trial wave function is to vary the mass m in $h(m)$. Trials at $m = -0.05$ and -0.1 did not produce any dramatic changes. This is an obvious place for future work.

The planar action with hypercubic fat links shows nearly identical behavior. Both these actions need about fifty per cent more inner CG steps than the (0.45,10) planar action but are still a large improvement over the Wilson overlap—and one could still run them on work stations. They might be useful in contexts where a very fat link might have undesirable properties.

Finally, the Gaussian action with (0.45,10) APE blocking requires about half as many inner CG calls as the (0.45,10) planar action, typically 3000-5000. However, this action requires about 2.5 times as much storage and 2.5 times as much CPU time per inner CG step as the planar overlap action, since its couplings span a hypercube (80 neighbors rather than 24). Thus it does not seem to be competitive with the planar action without clever coding [23].

The (0.45,10) fat clover action, with $C_{SW} = 1$, was also briefly investigated as a kernel for the overlap. It seemed to need 100,000-300,000 inner CG steps in the fiducial calculation, in 2500 to 8000 Rayleigh iterations, beginning from $m = 0$ trial wave functions. (There is an extreme variation in these numbers from lattice to lattice tested.) Eigenmodes of the fat link clover action are apparently quite different from those of the corresponding overlap action. The number of inner CG steps reflects the greater conditioning number of $h(-r_0)$.

These tests are certainly incomplete. It may be that I have simply written a very inefficient Wilson overlap, though most of the code for the two methods is common. My Wilson overlap is so expensive that it cannot be tuned, whereas

it is easy to test variations of the planar overlap. The “design philosophy” of beginning with an action which is close to an overlap action certainly produces actions which are inexpensive enough to run on small computers (work stations in my case).

IV. THE QUARK CONDENSATE IN FINITE VOLUME

In order to do a little physics in what is otherwise a paper about technique, I present a calculation of the chiral condensate in the quenched approximation with the (0.45,10) fattened planar overlap action. The method is (almost) exactly that of the pioneering calculation of Hernandez and Jansen and Lellouch [11]: one computes the condensate in background gauge field configurations of fixed topology labeled by winding number ν [24]. The condensate is

$$\Sigma_\nu = m\Sigma^2V(I_\nu(m\Sigma V)K_\nu(m\Sigma V) + I_{\nu+1}(m\Sigma V)K_{\nu-1}(m\Sigma V)) + \frac{\nu}{mV} \quad (8)$$

where m is the quark mass, V is the volume, and Σ is the infinite volume condensate. $I_\nu(z)$ and $K_\nu(z)$ are modified Bessel functions. In practice, one computes Σ “without topology,” by working in the chiral sector which has no zero eigenmodes (and doubling the result). Then one measures $\Sigma_\nu - \nu/(mV)$. In practice, the lattice number needs an additive renormalization: in zero mass, this is removed by the replacement of D^{-1} by $\tilde{D}^{-1}$ as in Eq. 5. There can also be contribution which vary with the quark mass, so the lattice number will need to be fit to

$$\langle \bar{\psi}\psi \rangle_{sub} = \frac{1}{V} \sum_x \tilde{D}_{x,x} = (\Sigma_\nu - \nu/(mV)) + Cm + \dots \quad (9)$$

One small difference between my calculation and that of Ref. [11] is in the normalization of the propagator: they do not have the prefactor $1/(1 - \frac{m}{2r_0})$ of Eq. 5. It is hard to believe that the condensate should not be an odd function of the symmetry breaking term (the quark mass), and the data does not show anything but a simple linear dependence on the mass.

Finally, Σ is scheme-dependent, and an overall lattice-to-continuum regulator needs to be computed.

Warned by the results of Ref. [11], I restricted my calculation to the $\nu = \pm 1$ sector. I generated a set of 40 lattices at each of three volumes 8^4 , 10^4 , and 12^4 at $\beta = 5.9$. I filtered them to find candidate $\nu = \pm 1$ configurations using a pure gauge measurement of topological charge [25], which I had previously calibrated against a set of 10^8 lattices. I checked that these lattices in fact had $\nu = \pm 1$ during the calculation of the condensate; only one lattice failed this test. This obviously leaves out configurations which the fermion observable identifies as carrying topological charge, but the gauge observable does not, but the alternative is to process every configuration through the overlap program. I was left with a set of 10^8 , $13 \cdot 10^4$, and $9 \cdot 12^4$ lattices on which I computed $\langle \bar{\psi}\psi \rangle$. I computed propagators at five quark masses, $am = 0.001, 0.002, 0.005, 0.01$, and 0.02 (again using a multi-mass Conjugate Gradient solver). I used twelve random sources per lattice (though I blocked data together before averaging). This needed about 40, 70, and 125 steps at the three volumes, at about 20-30 inner CG's per outer step, to reach a fractional squared residue of 10^{-12} . As a check, I also calculated Σ from the GMOR relation $\sum_x \langle \pi(x)\pi(0) \rangle = \Sigma/m$; as expected in the overlap it reproduced the direct calculation of Σ/m to within 10^{-7} (see Ref. [7]).

My lattice results are shown in Fig. 9.




FIG. 9. The quark condensate in finite volumes (divided by the quark mass). From the bottom, the volumes are 8^4 , 10^4 , and 12^4 . The curves are fits to Eq. 10.

The data at each volume are quite correlated. A single-elimination jackknife fit to

$$\langle \bar{\psi} \psi \rangle_{sub} = m \Sigma^2 V (I_1(m \Sigma V) K_1(m \Sigma V) + I_2(m \Sigma V) K_0(m \Sigma V)) + C m \quad (10)$$

gives (re-inserting the lattice spacing) $\Sigma a^3 = 0.00394(16)$, $C = 0.304(8)$. The lattice number is about two standard deviations higher than the number reported by Ref. [11] at $\beta = 5.85$ of $\Sigma a^3 = .0032(4)$. We can sharpen this disagreement by trading the lattice spacing a for the Sommer radius r_0 , using the interpolating formula of Ref. [26]: $\Sigma r_0^3 = 0.215(27)$ for Ref. [11], $\Sigma r_0^3 = 0.354(14)$ here.

A real comparison requires computing the Z factor. I have not done that yet, but I can make a heuristic attempt at the calculation, by exploiting the fact that perturbation theory for fat link actions becomes simple in the limit of large fattening. This argument is implicit in the discussion of fat link perturbation theory in Ref. [20]. Since it falls outside the main thrust of the paper, I relegate the discussion to an Appendix.

Taking r_0 to be 0.5 fm, $\Sigma_{\overline{MS}}(\mu = 2 \text{ GeV}) = 0.0244(10)(37) \text{ GeV}^3$. The two errors are from the lattice fit and from an assumed lattice spacing uncertainty of five per cent [27].

This calculation produces a number which disagrees badly with a calculation of the condensate using clover fermions and the GMOR relation, $\Sigma_{\overline{MS}}(\mu = 2 \text{ GeV}) = 0.0147(8)(16)(12) \text{ GeV}^3$ [28]. It is done at smaller lattice spacing and has completely different systematic uncertainties. Oddly enough, however, my result agrees well with an earlier analysis of data by Gupta and Bhattacharya: [30] $\Sigma_{\overline{MS}}(\mu = 2 \text{ GeV}) = 0.024(2)(2) \text{ GeV}^3$ (statistical and lattice spacing uncertainties).

Of course, all the potential weaknesses of this calculation, reported by Ref. [11] apply here, too: the lattices are small, and the lattice spacing is large. And it is a quenched calculation.

All of the computations of Σ were done over about a month of running on a set of four or five 450 Mhz Pentium II work stations.

V. CONCLUSIONS

None of the results shown here are particularly surprising, but that does not mean that they might be totally devoid of interest. A good approximate overlap action is easier to convert into an exact overlap, than any random action like the Wilson action. Crucial ingredients seem to be some kind of gauge connection which suppresses dislocations, a free action which “resembles” a free field overlap action, and the use of as much information from the trial $h(m)$ as possible to begin the calculation of the overlap action. One should be able to do better.

ACKNOWLEDGEMENTS

I am deeply indebted to Urs Heller for many conversations about the overlap action, to Robert Edwards for a table of Remes coefficients, to Kostas Orginos for a copy of his eigenvalue code, to Anna Hasenfratz for discussion about blocking, and to Archie Paulson for help coding. This work was supported by the U. S. Department of Energy.

APPENDIX A: Z-FACTORS FOR VERY FAT ACTIONS

The evaluation of the Z-factor is straightforward. I will use many of the techniques of Ref. [29], but collect them here for completeness. We expect $\Sigma(\mu)_{\overline{MS}} = Z(a\mu)\Sigma(a)_{latt}$ with $Z_S = 1/Z_m$ and Z_m is the quark mass renormalization factor. In perturbation theory, $Z_m = 1 + bz$, and

$$z = \Delta_{latt} - \Delta_{cont} = 6 \ln(1/(\mu a)) + x, \quad (A1)$$

where $b = \alpha_s(q^*)/(3\pi)$. I choose to use $\alpha_s = \alpha_{\overline{MS}}(q^*)$. I will use the plaquette to define a coupling $\alpha_V(3.41/a)$ ($=0.16054$ at $\beta = 5.9$), which will then be converted using two-loop perturbation theory to $\alpha_{\overline{MS}}(q^*)$, to give $\Sigma(\mu = 1/a)_{\overline{MS}}$. I will then run this result to $\mu = 2$ GeV, using the (inverse of the) two-loop running formula for the $\overline{MS}$ quark mass.

z is the difference between a lattice integral

$$\Delta_{latt} = \int_k I_{latt} F_N; \quad (A2)$$

and a continuum integral

$$\Delta_{cont} = \int_{k,c} I_{cont} \quad (A3)$$

I have factored the lattice integrand into a thin link piece I_{latt} and the form factor of the fat quark-gluon vertices F_N ; for APE-smeared links $F_N = (1 - \alpha \hat{q}^2/6)^{2N}$ where $\hat{q}^2 = \sum_\mu (4/a^2) \sin^2(k_\mu a/2)$. I am using the notation $\int_k = \int d^4k/(2\pi)^4$ over the hypercube, $-\pi/a < k_\mu < \pi/a$; $\int_{k,c} = \int d^4k/(2\pi)^4$. Introducing a gluon mass λ to regularize its IR divergence, $\Delta_{cont} = 5/2 - 3 \ln(\lambda^2 \mu^2)$ (in $\overline{MS}$). Adding and subtracting continuum-like terms to isolate the divergence in the lattice integral, we write

$$\begin{aligned} \Delta_{latt} &= \int_k (I_{latt} - 3 \frac{\theta(\pi^2 - k^2)}{k^4}) F_N - 3 \int_k \frac{\theta(\pi^2 - k^2)}{k^4} (1 - F_N) + 3 \int_k \frac{\theta(\pi^2 - k^2)}{k^4} \\ &= J - 3D + 3 \ln \frac{\pi^2}{a^2 \lambda^2}. \end{aligned} \quad (A4)$$

The first two terms (J and D) are IR and UV finite. Now the point is that D is a continuum-like integral, and all the dependence of Δ_{latt} on the lattice action is in the first term. But if the fattening is large, the form factor F_N , is nonzero only at tiny k , where it is unity. At that point, for all practical purposes, $I_{latt} \simeq 3/k^4$, and so we expect that J will be small. All the dependence of Z_m on the lattice action, other than its form factor, vanishes when we set

$J = 0$ in Eq. A4. q^* (defined according to the prescription of Lepage and Mackenzie [29]) is calculated by a similar procedure.

While I can't check this approximation for the overlap action, at (0.45,10) fattening, a complete calculation for clover fermions gives $x = -5.58$ and $q^* = 1.07$, while the $J = 0$ result is $x = -5.44$ and $q^* = 1.10$. A similar argument would predict that finite renormalization factors, such as the vector and axial current renormalizations, are unity, which is a good approximation to what is seen at large fattening. Of course, for small fattening, this approximation fails badly and a better (non-perturbative?) calculation is necessary. (For example, the (0.25,7) APE-blocked clover action has $x = -0.975$, $q^* = 2.88$; the approximate result is $x = -2.49$, $q^* = 1.59$.)

-
- [1] For complementary reviews, see F. Niedermayer, Nucl. Phys. Proc. Suppl. **73**, 105 (1999) [hep-lat/9810026] and H. Neuberger, Nucl. Phys. Proc. Suppl. **83-84**, 67 (2000) [hep-lat/9909042].
 - [2] H. Neuberger, Phys. Lett. B417, 141 (1998), Phys. rev. Lett. 81, 4060 (1998).
 - [3] P. Ginsparg and K. Wilson, Phys. Rev. D25, 2649 (1982).
 - [4] R. G. Edwards, U. M. Heller and R. Narayanan, Nucl. Phys. **B540**, 457 (1999) [hep-lat/9807017].
 - [5] P. Hernandez, K. Jansen and M. Lüscher, Nucl. Phys. **B552**, 363 (1999) [hep-lat/9808010].
 - [6] A. Borici, Phys. Lett. **B453**, 46 (1999) [hep-lat/9810064].
 - [7] R. G. Edwards, U. M. Heller and R. Narayanan, Phys. Rev. **D59**, 094510 (1999) [hep-lat/9811030].
 - [8] R. G. Edwards, U. M. Heller, J. Kiskis and R. Narayanan, Phys. Rev. **D61**, 074504 (2000).
 - [9] R. G. Edwards, U. M. Heller and R. Narayanan, Parallel Computing 25, 1395 (1999). hep-lat/9905028.
 - [10] R. G. Edwards, U. M. Heller, J. Kiskis and R. Narayanan, Phys. Rev. Lett. **82**, 4188 (1999)
 - [11] P. Hernandez, K. Jansen and L. Lellouch, Phys. Lett. **B469**, 198 (1999) [hep-lat/9907022].
 - [12] S. J. Dong, F. X. Lee, K. F. Liu and J. B. Zhang, hep-lat/0006004.
 - [13] See P. Hasenfratz and F. Niedermayer, Nucl. Phys. **B414**, 785 (1994) [hep-lat/9308004]; P. Hasenfratz, Nucl. Phys. **B525**, 401 (1998) [hep-lat/9802007]; hep-lat/9803027, plus references therein.
 - [14] F. Farchioni, I. Hip and C. B. Lang, Phys. Lett. **B443**, 214 (1998) [hep-lat/9809016]. F. Farchioni, C. B. Lang and M. Wohlgenannt, Phys. Lett. **B433**, 377 (1998) [hep-lat/9804012]; F. Farchioni, I. Hip, C. B. Lang and M. Wohlgenannt, Nucl. Phys. **B549**, 364 (1999) [hep-lat/9812018]; W. Bietenholz and I. Hip, Nucl. Phys. **B570**, 423 (2000) [hep-lat/9902019].
 - [15] T. DeGrand [MILC Collaboration], Phys. Rev. **D58**, 094503 (1998) [hep-lat/9802012]; T. DeGrand, A. Hasenfratz, and T. Kovacs, Nucl. Phys. B547, 259 (1999).
 - [16] T. DeGrand [MILC collaboration], Phys. Rev. **D60**, 094501 (1999) [hep-lat/9903006].
 - [17] : M. Albanese *et al.* [APE Collaboration], Phys. Lett. **B192**, 163 (1987); M. Falcioni, M. L. Paciello, G. Parisi and B. Taglienti, Nucl. Phys. **B251** (1985) 624.
 - [18] A. Hasenfratz, private communication.
 - [19] Cf. B. Jegerlehner, Nucl. Phys. (Proc. Suppl.) 63, 958 (1998), hep-lat/9612014; A. Frommer, B. Nockel, S. Güsken, T. Lippert and K. Schilling, Int. J. Mod. Phys. C6, 627 (1995) [hep-lat/9504020].
 - [20] C. Bernard and T. DeGrand, Nucl. Phys. Proc. Suppl. **83-84**, 845 (2000) [hep-lat/9909083].
 - [21] See B. Bunk, K. Jansen, M. Lüscher, and H. Simma, unpublished DESY report (1994); T. Kalkreuter and H. Simma, Comp. Phys. Comm. 93, 33 (1996).
 - [22] See, for example, the discussion in P. M. Vranas, hep-lat/0001006.
 - [23] P. Hasenfratz, S. Hauswirth, K. Holland, T. Jorg, F. Niedermayer and U. Wenger, hep-lat/0003013.
 - [24] J. C. Osborn, D. Toublan and J. J. Verbaarschot, Nucl. Phys. **B540**, 317 (1999) [hep-th/9806110]; P. H. Damgaard, J. C. Osborn, D. Toublan and J. J. Verbaarschot, Nucl. Phys. **B547**, 305 (1999) [hep-th/9811212].
 - [25] A. Hasenfratz and C. Nieter, Phys. Lett. **B439**, 366 (1998) [hep-lat/9806026].
 - [26] M. Guagnelli, R. Sommer and H. Wittig [ALPHA collaboration], Nucl. Phys. **B535**, 389 (1998) [hep-lat/9806005].
 - [27] A. X. El-Khadra, G. Hockney, A. S. Kronfeld and P. B. Mackenzie, Phys. Rev. Lett. **69**, 729 (1992), quote $1/a = 1.78(9)$ GeV, for a five per cent uncertainty, from the S-P splitting of charmonium.
 - [28] L. Giusti, F. Rapuano, M. Talevi and A. Vladikas, Nucl. Phys. **B538**, 249 (1999) [hep-lat/9807014].
 - [29] G. P. Lepage and P. B. Mackenzie, Phys. Rev. **D48**, 2250 (1993) [hep-lat/9209022].
 - [30] R. Gupta and T. Bhattacharya, Phys. Rev. **D55**, 7203 (1997) [hep-lat/9605039].